

INTRINSIC THERMODYNAMICAL TIME-SCALES OF THE ATMOSPHERE-OCEAN-CRYOSPHERE CLIMATE SYSTEM

Peter C. Chu

Department of Oceanography
Naval Postgraduate School
Monterey, CA 93943, USA

1. INTRODUCTION

Quantitative reconstruction of climatic parameters from both oceanic and continental records shows several time scales of the successive glacial-interglacial episodes that have characterized the Earth's climate about the past million years. For example, after analyzing data from the 2,083 m ice core recovered by the Soviet Antarctic Expeditions at Vostok (East Antarctic), Jouzel et al. (1982) found several peaks (25.3, 45.7, and 107.5 kyr) in the variation spectra of the Vostok isotope temperature, and related these peaks to the astronomical forcing, i.e., the obliquity of the Earth's axis (period at 41 kyr) and the precession of the equinox (periods at 23 and 19 kyr). Is the climatic variation purely caused by the external forcing? If so, the climate prediction would be relatively simple because it becomes the forecast of the Earth's orbit tilt and precession. In fact, the climate system is much more complicated. The climatic variation is caused by both external and internal sources (Fig.1).




Fig.1 A simple model of the climate system illustrating internal and external sources of climatic variation (after Imbrie, 1982).

The internal forcings are processes and mechanisms within the Earth's system, giving rise to self-excited climate fluctuations. The temperature-albedo feedback mechanism associated with long and short wave radiation fluxes at the Earth's surface are given considerable attention in glacial and climate models (e.g., Kallen et al., 1979; Oerlemans and van der Veen, 1984). However, not until recently (Chu, 1990), the role of hydrological cycle on the climatic variation on glacial and interglacial time-scales is given less attention although it is realized that ice freeze/melt and evaporation/precipitation are contributors to the change of air and ocean temperatures. Taking hydrological cycle into consideration, several intrinsic time scales of the atmosphere-ocean-cryosphere climate system will be discussed in this paper.

2. ATMOSPHERE-OCEAN-CRYOSPHERE CLIMATE SYSTEM

The total surface, ice covered, and open ocean areas are set to be 1, N_i , and $1 - N_i$, respectively (Fig.2). The interactions among the atmosphere, ocean, and ice is proposed by Chu (1990) as follows. From the global point of view, ice advance isolates the ocean from the atmosphere and reduces the heat loss from the ocean surface, which in turn decreases the air temperature and increases the ocean temperature. The air and ocean feed back into the ice through two different kinds of mechanisms (Fig.3): (1) thermodynamical feedback; the in-

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1993		2. REPORT TYPE		3. DATES COVERED 00-00-1993 to 00-00-1993	
4. TITLE AND SUBTITLE Intrinsic Thermodynamical Time-Scales of the Atmosphere-Ocean-Cryosphere Climate System				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Postgraduate School, Department of Oceanography, Monterey, CA, 93943				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Fourth Conference on Global Change, American Meteorological Society, 433-438					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

crease of ocean temperature melts the ice and causes the ice retreat (negative feedback between ice and ocean), however, the decrease of air temperature causes freezing and makes the ice further advance (positive feedback between ice and air); (2) mechanical feedback; the modification of the atmospheric and oceanic temperatures varies the surface evaporation rate, which in turn changes the sea-level height and the ice accumulation rate. Both effects lead to a further change of the ice flow.




Fig.2 Physical processes in the coupled atmosphere, ocean, and cryosphere climate system (after Chu, 1990).




Fig.3 Feedback mechanisms among air, ice, and ocean.

2.1 Atmosphere

The atmosphere is heated (or cooled) by radiation, and heat fluxes (including latent heat flux) from the air-ocean and air-ice interfaces, i.e.,

$$\frac{dT_a}{dt} = \frac{1}{\rho_a c_{pa} H_a} [-N_i Q_{ai} + (1 - N_i) Q_{wa} + R_a] \quad (1a)$$

where T_a is the air temperature, c_{pa} is the air specific heat, Q_{ai} , Q_{wa} are, respectively, the heat fluxes across air-ice and air-ocean interfaces, R_a is the net radiation absorbed by the atmosphere, and ρ_a and H_a are the characteristic values of atmospheric density and thickness.

In the present study, it is assumed that the water in the atmosphere is always in the vapor form. As soon as condensation happens, the condensed water droplets are assumed to be precipitation right away. Therefore, the time rate change of specific humidity q_a is caused by the excessive surface evaporation over precipitation and ice accumulation

$$\frac{dq_a}{dt} = \frac{1}{\rho_a H_a} [(1 - N_i) \rho_w (E - Pr) - N_i \rho_i c_s] \quad (1b)$$

where E , Pr , and c_s are rates of ocean surface evaporation, precipitation, and ice accumulation. ρ_w and ρ_i are the characteristic values of ocean and ice densities.

2.2 Ocean

Similar to the atmosphere, the ocean thermodynamics can be depicted by

$$\frac{dT_w}{dt} = -\frac{1}{\rho_w c_{pw} H_w} [N_i Q_{wi} + (1 - N_i) Q_{wa} + R_w] \quad (2)$$

where T_w is the ocean temperature, c_{pw} is the ocean specific heat, Q_{wi} is the heat fluxes across ice-ocean interfaces, R_w is the net radiation absorbed by the ocean, and H_w is the ocean thickness.

From a thermodynamical point of view, the ocean thickness is reduced by the excessive surface evaporation rate E over precipitation rate Pr , and by the ice edge freezing rate F ,

$$\frac{\partial}{\partial t} [(1 - N_i)\rho_w H_w] = -(1 - N_i)\rho_w(E - Pr) - (1 - N_i)\rho_i \Lambda(F)F - N_i \rho_i \Lambda(-F)F \quad (3)$$

where Λ is the Heaviside function, i.e., $\Lambda(x) = 1$, as $x \geq 0$, and $\Lambda(x) = 0$, as $x < 0$. $F > 0$, is taken as ice freezing; and $F < 0$, is taken as ice melting.

2.3 Ice-shelf

The ice is assumed to be isotropic with thickness H_i (Fig.2), and the ice-spreading rate is computed by (Weertman, 1957)

$$\langle \epsilon_h \rangle = A \left[\frac{1}{4} \rho_i g H_i \right]^n \quad (4)$$

where H_i is the height between the ice top and the sea-level (Fig.2). Values of n vary from about 1.5 to 4.2 with a mean of about 3, and for randomly oriented polycrystalline ice at -10°C and $n=3$, a value $A = 3 \times 10^{-8} \text{yr}^{-1} \text{kPa}^{-3}$ is reasonable for ice stress. The angle brackets denote values averaged over ice thickness.

The change of ice coverage N_i is due to ice-spreading and ice freezing/melting:

$$\frac{\partial N_i}{\partial t} = N_i \langle \epsilon_h \rangle + \frac{(1 - N_i)\Lambda(F)F}{H_i} + \frac{N_i \Lambda(-F)F}{H_i} \quad (5)$$

Following Killworth (1979), the heat flux is accomplished through an exchange coefficient K and is of the form $K(T - T_f)$, where T_f represents the freezing point of the sea water. The freezing rate is therefore computed by

$$F = -\frac{1}{\rho_i L_i} [\rho_w c_{pw} K_{wi}(T_w - T_f) + \rho_a c_{pa} K_{ai}(T_a - T_f)] \quad (6)$$

where K_{wi} , K_{ai} are the heat exchange coefficients for water-ice and air-ice, respectively. L_i is the latent heat of ice. If T_w , T_a are greater than T_f , F is taken as a negative value, which indicates ice melting.

Based on the continuity, Shumskiy (1965) proposed a method to determine the change of ice thickness by comparing the accumulation rate with horizontal strain-rates. Neglecting nonlinear inertial terms the time rate change of ice thickness becomes

$$\frac{\partial H_i}{\partial t} = c_s - H_i \langle \epsilon_h \rangle \quad (7)$$

The atmosphere-ocean-cryosphere climate system is depicted by (1), (2), (3), (5), and (7).

2.4 Interfacial Heat Fluxes

The heat fluxes across the air-ocean, air-ice, ice-ocean interfaces change the thermal features of each component of the atmosphere-ocean-cryosphere climate system. There are many different methods to compute these fluxes, among them, the simplest form was proposed by Killworth (1979):

$$\begin{aligned} Q_{ai} &= \rho_a c_{pa} K_{ai}(T_a - T_i) \\ Q_{wa} &= \rho_a c_{pa} K_{wa}(T_w - T_a) \\ Q_{wi} &= \rho_w c_{pw} K_{wi}(T_w - T_i) \end{aligned} \quad (8)$$

3. ROLE OF THE HYDROLOGICAL CYCLE

Summation of (1b), (3), (5) multiplied by H_i , and (7) multiplied by N_i leads to

$$\frac{d}{dt} [(1 - N_i)\rho_w H_w + \rho_i H_i N_i + \rho_a H_a q_a] = 0 \quad (9)$$

which shows the conservation of the water mass in the Earth system.

The water vapor evaporated from the global ocean can be estimated by

$$E = C_D |V_a| \frac{\rho_a}{\rho_w} [q_s(T_w) - q_a] \quad (10)$$

where C_D is the drag coefficient, V_a is the surface wind, q_s is the saturated mixing ratio, and q_a is the mixing ratio near the ocean surface. If the Earth surface is assumed to be covered by ocean and ice only, the water vapor evaporated from the ocean surface is transported by the atmospheric circulation, condensed and precipitated. The precipitation over ice leads to ice accumulation, therefore, if the total water vapor evaporated from the ocean surface is balanced by the precipitation over ocean (Pr) and the ice accumulation rate over ice (c_i), the relationship among surface evaporation rate, precipitation rate (over ocean), and ice accumulation rate should be

$$(1 - N_i)\rho_w(E - Pr) = N_i\rho_i c_s \quad (11)$$

4. TIME SCALES OF THE CLIMATE SYSTEM

In explaining the climatic variability, one may first ask: is climatic variability caused by external or internal forcings? The external forcings are agencies outside the Earth system, such as astronomically caused changes in the Earth's orbital parameters. In the atmosphere-ocean-cryosphere climate system

discussed in this paper, the external forcings mainly change the radiation absorbed by atmosphere (R_a) and ocean (R_w).

On the other hand, the internal forcings are processes and mechanisms within the Earth system, giving rise to self-excited climatic fluctuations. Substitution of (8) into (1) and (2), leads to thermal relaxation time scales for the atmosphere

$$\tau_a = \left[\frac{K_{at}}{H_a} N_i + \frac{K_{wa}}{H_a} (1 - N_i) \right]^{-1} \quad (12)$$

and for the ocean

$$\tau_w = \left[\frac{K_{wi}}{H_w} N_i + \frac{\rho_a c_{pa} K_{wa}}{\rho_w c_{pw} H_a} (1 - N_i) \right]^{-1} \quad (13)$$

The ice-spreading time scale can be defined from (5),

$$\tau_s = (\langle \epsilon_h \rangle)^{-1} \quad (14)$$

Horizontal ice growth (or ice edge freezing/melting) time scale is obtained from (5):

$$\tau_f = \frac{H_i}{F[\Lambda(F) - \Lambda(-F)]}$$

The perturbed ice accumulation c'_i is caused by the perturbed evaporation, precipitation, and the ice advancement:

$$c'_s = \frac{(1 - N_i)\rho_w}{N_i\rho_i} (E' - Pr') - [\rho_w(E - Pr) + \rho_i c_s] N'_i \quad (15)$$

The perturbations of evaporation and precipitation rates E' and Pr' vary on a much shorter time scale than the time-scale for ice advance/retreat. Another time scale for the ice growth in vertical direction can be defined by

$$\tau_t = \frac{\rho_i H_i}{[\rho_w(E - Pr) + \rho_i c_s]} \quad (16)$$

Here the variables without " ' " means the basic states. By the use of the values of the parameters listed in Table 1, these time scales are estimated as:

$$\begin{aligned}\tau_a &\approx 11.6d, \quad \tau_w \approx 66 \text{ yr}, \\ \tau_i &\sim 400 \text{ yr}, \quad \tau_s \sim 3000 \text{ yr}\end{aligned}$$

Since the atmospheric and oceanic thermal relaxation times scales are so much shorter than the ice-spreading time scale, the atmospheric and oceanic temperature perturbations T'_a and T'_w almost "instantaneously" follow the ice advance/retreat processes in the atmosphere-ocean-cryosphere climate system. From (1) and (2), we may obtain

$$T'_a = -A_a \tau_a N'_i, \quad T'_w = A_w \tau_w N'_i \quad (17)$$

where

$$A_a \equiv (Q_{wa} + Q_{ai})/(\rho_a c_{pa} H_a), \quad A_w \equiv Q_{wa}/(\rho_w c_{pw} H_w)$$

Equation (17) clearly shows the cooling of the atmosphere ($T'_a < 0$) and warming of the ocean ($T'_w > 0$) due to the ice advance. The perturbed ice freezing rate becomes

$$F' = -H_i(\alpha_w A_w \tau_w - \alpha_a A_a \tau_a) N'_i \quad (18)$$

where

$$\alpha_a \equiv \rho_a c_{pa} K_{ai}/(\rho_i L_i H_i), \quad \alpha_w \equiv \rho_w c_{pw} K_{wi}/(\rho_i L_i H_i)$$

The perturbed ice spreading rate is computed by

$$e'_h = n \tau_s^{-1} \frac{(h'_i - h'_w)}{H_i} \quad (19)$$

6. LOW FREQUENCY MODES GENERATED BY THE FEEDBACK MECHANISM

All the time scales presented in the last section are shorter than the time scales of the successive glacial-interglacial episodes (10-100 kyr) shown in Fig.1. If we consider the atmosphere-ocean-cryosphere as a one system,

the low frequency modes will be generated. If the coupled system perturbed from its equilibrium states, after mathematical manipulation, the perturbation equations are

$$\begin{aligned}\left[\frac{\partial}{\partial t} + (1 - \sigma) \tau_f^{-1} + N_i \mu_1 \mu_3 \tau_s^{-1} \right] N'_i \\ = \mu_2 N_i \tau_s^{-1} \frac{h'_i}{H_i}\end{aligned} \quad (20)$$

$$\begin{aligned}\left[\frac{\partial}{\partial t} + (1 + \mu_2) \tau_s^{-1} \right] \frac{h'_i}{H_i} \\ = (\mu_1 \mu_3 \tau_s^{-1} - N_i^{-1} \tau_i^{-1}) N'_i\end{aligned} \quad (21)$$

where

$$\mu_1 \equiv n \frac{\rho_w H_w - \rho_i H_i}{\rho_w H_w (1 - N_i)} \approx 2.431$$

$$\mu_2 \equiv n \left[1 + \frac{N_i \rho_i}{(1 - N_i) \rho_w} \right] \approx 3.66$$

$$\mu_3 \equiv \frac{H_w}{H_i} \approx 2.5$$

and

$$\sigma \equiv \tau_f (\alpha_a A_a \tau_a - \alpha_w A_w \tau_w)$$

The general solutions of equations (20) and (21) have the following forms:

$$(h'_i/H_i, N'_i) \sim e^{-i\omega t} \quad (22)$$

Substitution of (22) into (20) and (21) leads to a second order algebraic equation for ω

$$\begin{aligned}\omega^2 + i\omega[(1 - \sigma) \tau_f^{-1} + (1 + \mu_2 + N_i \mu_1 \mu_3) \tau_s^{-1}] - \\ [(1 - \sigma)(1 + \mu_2) \tau_s^{-1} \tau_f^{-1} + \mu_1 \mu_3 N_i \tau_s^{-2} + \mu_2 \tau_s^{-1} \tau_i^{-1}] = 0\end{aligned} \quad (23)$$

The frequency here is a complex number, i.e., $\omega = \omega_r + i\omega_i$, ω_i is a growth rate, and ω_r is the frequency.

7. FREQUENCIES FOR NEUTRAL STABILITY MODES

The modes with zero growth rate ($\omega_i = 0$) are called neutrally stable modes. The solution of (23) with $\omega_i = 0$ is

$$\omega_r = \tau_s^{-1} \{ \mu_2 v - [(1 + \mu_2)^2 + N_i \mu_1 \mu_2 \mu_3] \}^{1/2} \quad (24)$$

where

$$v \equiv \frac{\tau_s}{\tau_i}$$

The time scale for the atmosphere-ocean-cryosphere climate system depends on the ratio of the two shorter time scales: τ_s and τ_i . Fig. 4 shows the dependence of periods ($2\pi/\omega_r$) of the neutrally stable modes on the ratio v . When $v = 7.2$, the period is 43.3 kyr, which indicates that the feedback mechanisms among atmosphere, ocean, and cryosphere can also generate climate change on 10-100 kyr time scales.




Fig. 4 Periods versus τ_s/τ_i for the marginal stable modes.

DISCUSSION

The conventional theories refer the climate change on 10-100 kyr time scales to the concentration of variance near the Earth's orbit tilt and precession frequencies (Jouzel et al., 1987). However, the conceptual atmosphere-ocean-cryosphere model described in this paper shows a possible positive/negative feedback mechanism, induced by the hydrological cycle.

ACKNOWLEDGEMENTS

This research was sponsored by the National Science Foundation (DPP-9114161) and the Naval Postgraduate School.

REFERENCES

- Chu, P.C., 1990: Air-ice-ocean feedback mechanisms and ice oscillation on millennial time scales. *Ann. Glaciol.*, **14**, 28-31.
- Josberger, E.G., 1987: Bottom ablation and heat transfer coefficients from the 1983 Marginal Ice Zone Experiments. *J. Geophys. Res.*, **92**, 7012-7016.
- Jouzel, J., C. Lorius, J.R. Petit, C. Genthon, N.I. Barkov, V.M. Kotlyakov, and V.M. Petrov, 1987: Vostok ice core: a continuous isotope temperature record over the last climatic cycle (160,000 years). *Nature*, **329**, 403-407.
- Kallen, E., C. Crafoord, and M. Ghil, 1979: Free oscillations in a climate model with ice-sheet dynamics. *J. Atmos. Sci.*, **36**, 2292-2303.
- Killworth, P.D., 1979: On "chimney" formations in the ocean. *J. Phys. Oceanogr.*, **9**, 531-554.
- Oerlemans, J. and C.J. van der Veen, jr., 1984: *Ice Sheets and Climate*, Reidel Publishing Company, Dordrecht.
- Shumskiy, P.A., 1965: A study of ice in the Arctic Ocean. *Dokl. Akad. Nauk USSR*, **162**, 320-322.
- Weertman, J., 1957: Deformation of floating ice shelves. *J. Glaciol.*, **3**, 38-42.